

# The ornithologist Alfred Russel Wallace and the controversy surrounding the dinosaurian origin of birds

Nizar Ibrahim · Ulrich Kutschera

Received: 18 April 2013 / Accepted: 27 June 2013 / Published online: 24 August 2013  
© Springer-Verlag Berlin Heidelberg 2013

**Abstract** Over many years of his life, the British naturalist Alfred Russel Wallace (1823–1913) explored the tropical forests of Malaysia, collecting numerous specimens, including hundreds of birds, many of them new to science. Subsequently, Wallace published a series of papers on systematic ornithology, and discovered a new species on top of a volcano on Ternate, where he wrote, in 1858, his famous essay on natural selection. Based on this hands-on experience, and an analysis of an *Archaeopteryx* fossil, Wallace suggested that birds may have descended from dinosaurian ancestors. Here, we describe the “dinosaur-bird hypothesis” that originated with the work of Thomas H. Huxley (1825–1895). We present the strong evidence linking theropod dinosaurs to birds, and briefly outline the long and ongoing controversy around this concept. Dinosaurs preserving plumage, nesting sites and trace fossils provide overwhelming evidence for the dinosaurian origin of birds. Based on these recent findings of paleontological research, we conclude that extant birds indeed descended, with some modifications, from small, Mesozoic theropod dinosaurs. In the light of Wallace’s view of bird origins, we critically evaluate recent opposing views to this idea, including Ernst Mayr’s (1904–2005) arguments

against the “dinosaur-bird hypothesis”, and document that this famous ornithologist was not correct in his assessment of this important aspect of vertebrate evolution.

**Keywords** Alfred Russel Wallace · Evolution · Dinosaurs · Birds · *Archaeopteryx* · Ernst Mayr

## Introduction

Seven years after returning to his home country of England, Alfred Russel Wallace (1823–1913) published his most commercially successful book, *The Malay Archipelago* (Wallace 1869). During his 8-year-long excursions (1854–1862) into these tropical regions of planet Earth, the British naturalist independently discovered the principle of natural selection in populations of wild animals (Wallace 1858) and proposed a biogeographical concept that was later called “Wallace line”. In addition to these contributions to evolutionary biology and biogeography, he was also an avid collector and discovered/described many new species of birds.

In one of his original papers dealing with systematic ornithology, Wallace (1864) provided information on the distribution and habits of parrots of the Malayan region, with the description of two new species. In another research paper on this subject, Wallace (1865) characterized 21 species of birds from this area, and pointed out that he had, at that time, collected a total of 212 new species of birds in the islands of the Malay Archipelago.

Another of his many original contributions to ornithology dealt with the raptorial birds of the Malay Archipelago. The author remarked, “With the exception that Vultures are entirely absent, birds of prey are tolerably plentiful in the Archipelago” (Wallace 1868, p. 2). Interestingly, the British ornithologist attributed the occurrence of many

---

This article is a contribution to the Special issue Alfred Russel Wallace (1823–1913): The man in the shadow of Charles Darwin—Guest Editors U. Kutschera, U. Hossfeld.

---

N. Ibrahim  
Department of Organismal Biology and Anatomy,  
University of Chicago, 1027 East 57th Street, Chicago,  
IL 60637, USA  
e-mail: nibrahim@uchicago.edu

U. Kutschera (✉)  
Institute of Biology, University of Kassel, Heinrich-Plett-Str. 40,  
34132 Kassel, Germany  
e-mail: kut@uni-kassel.de

species to geographic separation of populations. Hence, Wallace foresaw not only Ernst Mayr's (1904–2005) biological species concept (Kutschera 2003, 2009), but also the principle of allopatric speciation, when he suggested that “This large number (of bird species) seems to be chiefly because of the breaking up of the district into a vast number of islands” (Wallace 1868, p. 2).

Finally, in his best- and long-seller *The Malay Archipelago*, Wallace (1869) devoted many pages to the distribution, behavior and taxonomy of different members of the Aves. His lucid descriptions of the exotic birds of paradise were one of the reasons for the tremendous success of this book, which is still in print today.

In this article, we discuss the opinions of Wallace concerning the evolutionary origin of birds, a class of vertebrates that he studied over many decades of his long life (Fig. 1) with reference to recent discoveries that have shed much needed light on the evolutionary history of this group of tetrapods.

### Alfred Russel Wallace and the origin of birds


Despite the fact that Wallace collected more than 8,000 specimens of birds and described many new species (Wallace 1869) during his travels in Southeast Asia, his writings on the evolutionary origin of birds are rather limited. His discussions of the evolutionary history of other groups are more extensive. In his popular book *Darwinism*, Wallace (1889) devoted chapter XIII to the “Geological evidences of evolution”. With reference to the science of

geology, the author pointed out that “The theory of evolution in the organic world necessarily implies that the forms of animals and plants have, broadly speaking, progressed from a more generalized to a more specialized structure, and from simpler to more complex forms” (Wallace 1889, p. 182). Wallace discussed the evolution of horses and other animal taxa, with reference to the corresponding work of the eminent British biologist Thomas Henry Huxley (1825–1895). In one sentence, Wallace (1889) also referred to the origin of birds. In his discussion of Huxley's research on fossil crocodiles, he remarked that “Dinosaurians (are) reptiles which in some respects approach birds” (Wallace 1889, p. 184). The author agreed with Huxley's opinion and, on a more general note, argued that the fossil finds of his time provide “evidence of evolution; the doctrine resting upon exactly as secure a foundation as did the Copernican theory of the motions of the heavenly bodies at the time of its promulgation” (Wallace 1889, p. 187).

In his popular book *The World of Life*, Wallace discussed several species of dinosaurs, and devoted one section to “The Earliest Birds” (Wallace 1910). He depicted and discussed the then famous “*Archaeopteryx macrura*”, figured the skull of *A. siemensii* (Fig. 2), and noted that “The very earliest known fossil bird is from the Upper Jurassic of Bavaria, and is beautifully preserved in the fine-grained beds of lithographic stone ... it is a true bird, notwithstanding its curiously elongated tail feathered on each side”. In the legend to his figure, he described this



**Fig. 1** Drawing of the “Ternate-bird” (*Rhipidura torrida*), discovered and described by Alfred R. Wallace on the summit of the volcano, 4,000 feet above the sea, on Ternate island (adapted from Wallace 1865)



**Fig. 2** Drawing of the “fossil lizard-tailed bird” (“*Archaeopteryx macrura*”—the London *Archaeopteryx*), and the skull of *Archeopteryx siemensii* (the Berlin specimen). (adapted from Wallace 1910)

fossil as a “Lizard-Tailed Bird” (Wallace 1889, pp. 213–215). He also explained why bird remains in the Mesozoic era are rare, but Wallace did not explicitly refer to the “dinobird-concept”, which was proposed several years before by his colleague Huxley, whom he treated with admiration and respect.

### The dino-bird hypothesis: origin and evolution of a concept

The dinosaurian origin of birds, originating as an idea in the 1860s with Huxley (1868a, b, 1870), is widely accepted today (Prum 2002; Chiappe 2004, 2007, 2009; Erickson et al. 2009) and remains a very active and dynamic area of research. The re-emergence of the view that dinosaurs are in fact bird-like creatures is relatively recent, but first mentions of the shared anatomical characteristics between birds and (non-avian) dinosaurs, date back much further. Early pioneers of archosaur comparative anatomy, such as the German anatomist Carl Gegenbaur (1826–1903) (Hossfeld et al. 2003), and the aforementioned zoologist Huxley, recognized these similarities (Wagner 1861; Chiappe 2007), which are most apparent in the skeletons of theropod dinosaurs. Thomas H. Huxley, today widely known as “Darwin’s bulldog”, was one of the most vociferous proponents of a bird-dinosaur link (Huxley 1868a, b, 1870).

While Huxley had little doubt about the dinosaurian origin of birds, some authors considered other scenarios,



**Fig. 3** Photograph of the Berlin *Archaeopteryx lithographica* (or *siemensii*; see Mayr et al. 2007). Picture courtesy of the Museum für Naturkunde, Berlin (Germany)



**Fig. 4** Reconstruction of *Archaeopteryx*. Artwork by Davide Bona-donna. *Archaeopteryx lithographica*-specimens typically represent animals roughly the size of a magpie, although some were larger, about the size of a raven (see Erickson et al. 2009)



**Fig. 5** Photograph of the Munich specimen of the small predatory dinosaur *Compsognathus longipes* Wagner (1861), which was interpreted as a bird-like reptile. The total body length of large *Compsognathus* exceeded 1 m (the Munich specimen is considerably shorter and represents a juvenile; see Therrien and Henderson 2007)

such as a thecodont origin as more likely (Heilmann 1926). These views became widespread following the publication of Heilmann’s influential book *The Origin of Birds*, aided to no small extent by the incredibly vibrant life illustrations Heilmann included in this work. By the early twentieth century, dinosaurs were no longer thought to be particularly relevant to bird origins, despite the existence of a truly astonishing fossil from Europe.

Because of the incompleteness of the fossil record, the world-famous “Urvogel” *Archaeopteryx*, from the Jurassic of Southern Germany (Bavaria), was for a long time the only key taxon and provided important insights into the morphology of a “transitional form”: feathers were clearly visible in this taxon, but so were toothed jaws, well-developed claws, and a long, dinosaurian tail (Figs. 3, 4). Huxley wrote about the similarities *Archaeopteryx* shared with small predatory dinosaurs, such as *Compsognathus* (Figs. 3, 4, 5), also from the Jurassic of Germany (Huxley 1868b). These two taxa in particular look remarkably similar in their skeletal anatomy and body shape (Figs. 2, 3, 4, 5).

Considering the huge scientific importance, beautiful preservation, and rarity of *Archaeopteryx* fossils, it was probably only a question of time before doubts about the authenticity of the fossil were voiced. Claims that *Archaeopteryx* is a forgery were quickly and thoroughly proven to be false (Charig et al. 1986), and several new specimens have been discovered (Mayr et al. 2007); they are all referred to *A. lithographica* by some authors (Houck et al. 1990; Senter and Robins 2003), while others distinguish two or more species (Mayr et al. 2007). For a long time, *Archaeopteryx* remained one of the very few physical pieces of evidence showing a mosaic of easily identifiable traditional “reptile” and “bird” characteristics.

Part of the fascination of the dinosaur-bird link undoubtedly has to do with the widespread, but superficial perception of a great difference in size and morphology between dinosaurs such as *Tyrannosaurus rex* and the vast majority of modern birds. In reality, many dinosaur skeletons show a striking similarity to the skeleton of a modern bird. This perception has sometimes hindered research exploring a possible relationship, and dinosaurs were generally thought to lack many typical avian features, such as a furcula, (proto) feathers, or an avian-style brooding behavior.

A pronounced shift in the perception of dinosaurs by the scientific community took place in the 1960s: away from a more “reptilian”, cold-blooded dinosaur (Colbert 1961), and towards a possibly endothermic, active, and more bird-

like “reptile” (Chiappe 2009). This view was strongly espoused by a small group of dinosaur paleontologists, among them John Ostrom (1928–2005) and Robert Bakker (Bakker 1968; Ostrom 1973). Ostrom’s discovery of *Deinonychus antirrhopus* in particular proved very important (Ostrom 1969). This predatory dinosaur is extremely bird-like in its anatomy (Fig. 6) and possibly even behavior, as suggested, among other things, by its appendicular anatomy (Fowler et al. 2011) and brooding behavior (Grellet-Tinner and Makovicky 2006). More recent finds have extended many of these features to other derived theropods, especially the group known as the maniraptoran dinosaurs (Norell et al. 1995).

### Fossil evidence from Asia and beyond

Today, after an ever-increasing rate of discovery of spectacular fossils, the number of shared anatomical characters between avian and non-avian dinosaurs is overwhelming. Numerous fossil theropod skeletons (e.g., Xu and Norell 2004; Norell and Xu 2005), nesting sites (e.g., Norell et al. 1995; Clark et al. 1999; Varricchio and Jackson 2004), ootaxa (e.g., Varricchio and Jackson 2004; Grellet-Tinner et al. 2006) and ichnofossils (e.g., Milner et al. 2009) document the strong evidence for a dinosaurian origin of birds (e.g., Sereno 1999; Chiappe and Witmer 2002). The finds are, for the most part, from Asia (China and Mongolia). They include dinosaurs preserved in an avian sleeping posture (Xu and Norell 2004), dinosaurs sitting on a clutch of eggs (Norell et al. 1995; Clark et al. 1999; Varricchio and Jackson 2004), adopting a bird-like resting pose (Milner et al. 2009) and outlines of fully feathered limb- or tail-sections (Xu et al. 2003; Chiappe 2004) (Fig. 7).

In terms of skeletal evidence, it is now well established that theropod dinosaurs share with birds a large number of key features (Fig. 8), including but not limited to: a bipedal posture, extensive pneumatization of the skeleton (O’Connor and Claessens 2005; Sereno et al. 2008), the presence of a furcula, swivel-like wrist joints (semi lunate

**Fig. 6** Reconstruction of a feathered *Deinonychus antirrhopus*. The morphology and distribution of the plumage is based on recent paleontological discoveries from Asia. Artwork by Davide Bonadonna. Total body length of *Deinonychus*: ca. 3.3 m



**Fig. 7** A representative *Microraptor zhaoianus* fossil showing body, wing, hind limb and tail feathers. Photograph courtesy of Mick Ellison (American Museum of Natural History). Scale bar 5 cm



carpal), L-shaped coracoids, and fast growth (Erickson et al. 2001). The large number of key discoveries cementing the role of predatory dinosaurs as bird ancestors led to intense debates about the origin of flight (Padian and de Ricqlès 2009). Clearly, some forms could not fly and used feathers for display and/or insulation, but others did have the anatomical features necessary for flight (Chiappe 2004).

Of particular interest in the evolution of flight are, in addition to *Archaeopteryx*, several new Asian taxa, including four-winged forms such as *Microraptor* (Fig. 7), a small maniraptoran from Asia (Xu et al. 2003; Li et al. 2012). These new specimens add important information to our understanding of leg feather morphology and evolution (Zheng et al. 2013). The precise sequence of events leading from flightless dinosaurs to full-powered bird flight is still debated and different models have been proposed (Chiappe 2009; Dececchi and Larsson 2011). It is clear, however, that plumage did initially evolve for purposes unrelated to flight.

### Frauds and unresolved questions

Unfortunately, research into the dinosaurian origin of birds has been temporarily overshadowed by examples of fraud, in particular, specimens that represent composite slabs, consisting of several taxa (Zhou et al. 2002). The most famous of these fabrications has been named “*Archaeoraptor*” and, because a series of unfortunate events meant that the forgery was not detected before initial description, this provided unnecessary “ammunition” to critics of the dinosaurian origin of birds, from sceptical ornithologists to US creationists. As is the case with other “Piltdown Man” style forgeries, this unfortunate episode should not detract from the fact that in the time elapsed since the “*Archaeoraptor*” story, countless new genuine and spectacular fossils of avian and non-avian dinosaurs have come to light, in addition to the ones already known. These new finds provide a spectacular fossil sequence, rich in diversity


and morphology and indicative of a complex evolutionary branching pattern.

Because of this complexity in theropod and bird evolution, some important details of taxonomic relationships remain unresolved, as competing hypotheses are currently not strong enough to rule each other out. This is relevant to the position of certain groups of theropod dinosaurs such as dromaeosaurs, troodontids, alvarezsaurids and oviraptorids for example—all of which have at one point or another been described as theropods that are particularly close to birds, and in some cases even as secondarily flightless birds (Perle et al. 1993; Chiappe et al. 1998; Sereno 2001; Hwang et al. 2002; Paul 2002; Witmer 2002).

More recently, evolutionary developmental biology (evo-devo) has also become an important area of debate, especially regarding digit homologies. This has traditionally been a divisive and complicated issue, but data from developmental biology now seem to show little incompatibility with the theropod origin of birds (Naish 2011). This is also supported by recent studies based on molecular evidence (Vargas and Fallon 2005).

Another topic discussed in avian evolution is what some researchers refer to as the “temporal paradox” (Feduccia 1999), which, in this case, describes the temporal discrepancy between the Cretaceous age of many of the new Asian dinosaurs showing protofeathers, feathers and wings, and the appearance of the earliest known bird, *Archaeopteryx*, which is reliably dated to the Late Jurassic (~150 million years ago) (Fig. 3). This criticism has, however, little merit, since (a) maniraptoran dinosaurs are in fact known from remains that are early Jurassic in age and that consequently predate *Archaeopteryx* (e.g., Witmer 2002; Hu et al. 2009), (b) it has not been proposed that the Cretaceous forms are ancestral to *Archaeopteryx* or similar forms, and (c) all previously proposed alternatives are far less parsimonious (Brochu and Norell 2001; Witmer 2002).

The position of *Archaeopteryx* has also recently been questioned (Xu et al. 2011), because one analysis recovered the taxon outside of Aves. This result should, however, not be regarded as a major turning point. While



**Fig. 8** Evolution of the avian skeleton from theropod dinosaurs during the Mesozoic, with representative species. The upper boundary of the timeline is marked by the Cretaceous/Paleogene mass

extinction. Only one lineage (Euornithes, 9), leading to extant birds, survived the extinction event (adapted from Sereno 1999)

cladistic analyses are powerful and important tools, they can suffer from a lack of data and/or stability and results obviously also depend on definitions of groups (in this case Aves). Indeed, the phylogenetic position of *Archaeopteryx* has subsequently been reverted back to its “original” position (Lee and Worthy 2012), emphasizing the fragility

of some phylogenetic analyses, and the sometimes exaggerated importance given to them by some authors to the detriment of other approaches, such as classical comparative anatomy. This problem was the cause for several misidentifications in the past, including some concerning early birds (Naish et al. 2012; Buffetaut 2011).

## Ernst Mayr and ongoing opposition against the dino-bird hypothesis

Unlike Alfred R. Wallace (1889, 1910), some modern ornithologists have opposed the dinosaurian origin of birds. This view was also espoused by one of the most influential evolutionary biologists of the twentieth century, Ernst Mayr. This world-expert in ornithology discovered and described many species of birds (Haffer 2007). In his popular book *What Evolution Is*, Mayr (2001), like some other ornithologists, appears to have underestimated, and possibly misunderstood some of the evidence linking the-ropod dinosaurs to birds.

Mayr (2001) argued that there are two distinct, plausible theories on the origin of birds, and erroneously states that the theory of the dinosaurian ancestry of birds suggests an origin from Cretaceous theropods. This is incorrect for the reasons explained above (see temporal paradox). The other theory Mayr refers to is the poorly supported thecodont origin (Witmer 2002), which suggests that birds originated from another archosaur group in the Triassic (Martin 2004). Mayr (2001) lists several, now largely discredited, lines of refutations against a dinosaurian origin of birds: differences in digit counts; differences in tooth morphology; the pectoral anatomy of late Cretaceous dinosaurs, and problems with the “ground up” model of avian flight. While a detailed refutation of the above arguments is beyond the scope of this paper and can be found elsewhere (e.g., Witmer 2002; Chiappe 2009), it is worth noting that some of them were surprisingly weak, even before the more recent discoveries described above made them untenable. To illustrate just one example, the argument that theropod teeth are more flattened than the simpler peg-like teeth of early birds (Mayr 2001) ignores the fact that theropod dentition, while often relatively similar, can also vary greatly in morphology, from conical, rounded teeth in spinosaurids (Dal Sasso et al. 2005), via the tiny, heterodont and numerous teeth of *Pelecanimimus* (Perez-Moreno et al. 1994), to the blade-shaped teeth of carcharodontosaurids (Serenio et al. 1996), to cite just a few examples. There is little doubt that in the face of such diversity, tooth morphology seems to be a poor choice for argument.

The view Mayr (2001) summarized in his popular book is in stark contrast to that expressed by the vast majority of vertebrate paleontologists, but emblematic for a relatively deep historical schism between the latter, and a substantial number of ornithologists (Feduccia 1999). This may be in some cases a consequence of a general lack of interest, and/or expertise, in the methods used in vertebrate paleontology. As the evidence for the dinosaurian origin of birds has continuously piled up, this divide has slowly been reduced, but it has not completely disappeared (Naish 2011). Based on currently available information and recent discoveries,

the prominent ornithologist Mayr (2001) was not correct in his assessment of this particular aspect of evolutionary biology.

## Conclusions

In two of his most influential books, Wallace (1889, 1910) discussed the evolutionary origin of birds. Based on the evidence available at that time, the ornithologist Wallace pointed out that extant birds may be the descendants of extinct dinosaurs, but he did not elaborate on this topic. This “dinobird-hypothesis” was proposed for the first time by the Victorian naturalist Thomas H. Huxley, who published several original papers on this topic (Huxley 1868a, b, 1870).

Today, it is widely accepted that birds, or “avian dinosaurs”, descended from, and in fact are, theropods that survived the Cretaceous/Paleogene mass extinction event 65 million years ago (Chiappe 2009; Kutschera 2013) (Fig. 8). Small predatory dinosaurs are the evolutionary ancestors of extant members of the Aves, a class of homeothermic vertebrates that comprises ca. 10,000 extant species (van Tuinen 2009). Like Wallace, Ernst Mayr was a remarkable ornithologist who discovered and described numerous species of birds (Haffer 2007). However, in his influential book *What Evolution Is*, Mayr (2001) summarized evidence against Huxley’s “dinobird hypothesis” in its “evolved” version (Feduccia 1999).

In this article, we have shown that Mayr’s writings do not reflect the current understanding of bird origins, and conclude that these feathered vertebrates are in fact a group of highly modified theropods that evolved during the Mesozoic from small, feathered dinosaur ancestors.

**Acknowledgments** The authors would like to thank Carol Abraczinskas for formatting and improving the figures, and Carol Abraczinskas and Paul Sereno for providing Fig. 8. Mick Ellison (American Museum of Natural History) kindly provided the photograph used in Fig. 7. Davide Bonadonna provided the artwork shown in Figs. 4 and 6. For comments on this manuscript, we would like to thank Alex Dececchi, Simone Maganuco and the reviewer, Daniela Schwarz-Wings.

## References

- Bakker RT (1968) The superiority of dinosaurs. *Discovery* 3:11–22
- Brochu CA, Norell MA (2001) Time and trees: A quantitative assessment of temporal congruence in the bird origins debate. In: Gauthier J, Gall LF (eds) *New perspectives on the origin and early evolution of birds*. Yale Peabody Museum, New Haven, pp 511–535
- Buffetaut E (2011) *Samrukia nessovi*, from the Late Cretaceous of Kazakhstan: a large pterosaur, not a giant bird. *Annales de Paléontologie* 97:133–138

- Charig AJ, Greenway F, Milner AC, Walker CA, Whybrow J (1986) *Archaeopteryx* is not a forgery. *Science* 232:622–626
- Chiappe LM (2004) The closest relatives of birds. *Ornitología Neotropical* 15:101–116
- Chiappe LM (2007) *Glorified dinosaurs: the origin and early evolution of birds*. John Wiley and Sons, New York
- Chiappe LM (2009) Downsized dinosaurs: the evolutionary transition to modern birds. *Evo Edu Outreach* 2:248–256
- Chiappe LM, Witmer LM (2002) *Mesozoic birds: above the heads of dinosaurs*. University of California Press, Berkeley
- Chiappe LM, Norell MA, Clark JM (1998) The skull of a relative of the stem-group bird *Mononykus*. *Nature* 392:275–278
- Clark JM, Norell MA, Chiappe LM (1999) An oviraptorid skeleton from the Late Cretaceous of Ukhaa Tolgod, Mongolia, preserved in an avian-like brooding position over an oviraptorid nest. *American Museum Novitates* 3265:1–36
- Colbert EH (1961) *Dinosaurs: Their discovery and their world*. WW Dutton, New York
- Dal Sasso C, Maganuco S, Buffetaut E, Mendez MA (2005) New information on the skull of the enigmatic theropod *Spinosaurus*, with remarks on its size and affinities. *J Vertebr Paleontol* 25:888–896
- Dececchi TA, Larsson HCE (2011) Assessing arboreal adaptations of bird antecedents: testing the ecological setting of the origin of the avian flight stroke. *PLoS ONE* 6:e22292
- Erickson GM, Curry-Rogers K, Yerby SA (2001) Dinosaurian growth patterns and rapid avian growth rates. *Nature* 412:429–433
- Erickson GM, Rauhut OWM, Zhou Z, Turner AH, Inouye BD, Hu D, Norell MA (2009) Was dinosaurian physiology inherited by birds? Reconciling slow growth in *Archaeopteryx*. *PLoS ONE* 4:e7390
- Feduccia A (1999) *The origin and evolution of birds*. 2nd edn. Yale University Press, New Haven
- Fowler DW, Freedman EA, Scanella JB, Kambic RE (2011) The predatory ecology of *Deinonychus* and the origin of flapping in birds. *PLoS ONE* 6:e28964
- Grellet-Tinner G, Makovicky P (2006) A possible egg of the dromaeosaur *Deinonychus antirrhopus*: phylogenetic and biological implications. *Can J Earth Sci* 43:705–719
- Grellet-Tinner G, Chiappe LM, Norell M, Bottjer D (2006) Dinosaur eggs and nesting behaviors: their paleobiological inferences. *Palaeogeogr Palaeoclimatol Palaeoecol* 232:294–321
- Haffer J (2007) *Ornithology, evolution, and philosophy. The life and science of Ernst Mayr 1904–2005*. Springer Verlag, Berlin/Heidelberg
- Heilmann G (1926) *The origin of birds*. HFG Witherby, London
- Hossfeld U, Olsson L, Breidbach O (2003) Editorial: Carl Gegenbaur (1826–1903) and his influence on the development of evolutionary morphology. *Theory Biosci* 122:105–108
- Houck MA, Gauthier JA, Strauss RE (1990) Allometric scaling in the earliest fossil bird, *Archaeopteryx lithographica*. *Science* 247:195–198
- Hu D, Hou L, Zhang L, Xu X (2009) A pre-*Archaeopteryx* troodontid theropod from China with long feathers on the metatarsus. *Nature* 461:640–643
- Huxley TH (1868a) Remarks upon *Archaeopteryx lithographica*. *Proc Zool Soc Lond* 16:243–248
- Huxley TH (1868b) On the animals which are most nearly intermediate between birds and reptiles. *Geol Mag* 5:357–365
- Huxley TH (1870) Further evidence of the affinity between the dinosaurian reptiles and birds. *Quart J Geol Soc* 26:12–31
- Hwang SH, Norell MA, Ji Q (2002) New specimens of *Microraptor zhaoianus* (Theropoda: Dromaeosauridae) from Northeastern China. *American Museum Novitates* 3381:1–44
- Kutschera U (2003) A comparative analysis of the Darwin—Wallace papers and the development of the concept of natural selection. *Theory Biosci* 122:343–359
- Kutschera U (2009) Symbiogenesis, natural selection, and the dynamic Earth. *Theory Biosci* 128:191–203
- Kutschera U (2013) Evolution. In: Maloy S, Hughes K (eds) *Brenner's Encyclopedia of genetics*, vol 2. Elsevier, New York, pp 541–544
- Lee MSY, Worthy TH (2012) Likelihood reinstates *Archaeopteryx* as a primitive bird. *Biol Lett* 8:299–303
- Li Q, Gao KQ, Meng Q, Clarke JA, Shawkey MD, D'Alba L, Pei R, Ellison M, Norell MA, Vinther J (2012) Reconstruction of *Microraptor* and the evolution of iridescent plumage. *Science* 335:1215–1219
- Martin LD (2004) A basal archosaurian origin for birds. *Acta Zool Sinica* 50:978–990
- Mayr E (2001) *What evolution is*. Basic Books, New York
- Mayr G, Phol B, Hartman S, Peters DS (2007) The tenth skeletal specimen of *Archaeopteryx*. *Zool J Linn Soc B* 149:97–116
- Milner ARC, Harris JD, Lockley MG, Kirkland JI, Matthews NA (2009) Bird-like anatomy, posture, and behavior revealed by an Early Jurassic theropod dinosaur resting trace. *PLoS ONE* 4:e4591
- Naish D (2011) *Glorified dinosaurs: the origin and early evolution of birds* (Book review). *Hist Biol* 23:435–438
- Naish D, Dyke G, Cau A, Escuillié F, Godefroit P (2012) A gigantic bird from the Upper Cretaceous of Central Asia. *Biol Lett* 8:97–100
- Norell MA, Xu X (2005) Feathered dinosaurs. *Annu Rev Earth Planet Sci* 33:277–299
- Norell MA, Clark JM, Chiappe LM, Dashzeveg D (1995) A nesting dinosaur. *Nature* 378:774–776
- O'Connor PM, Claessens LPAM (2005) Basic avian pulmonary design and flow-through ventilation in non-avian dinosaurs. *Nature* 436:253–256
- Ostrom JH (1969) Osteology of *Deinonychus antirrhopus*, an unusual theropod from the Lower Cretaceous of Montana. *Peabody Museum of Natural History Bulletin* 30:1–165
- Ostrom JH (1973) The ancestry of birds. *Nature* 242:136
- Padian K, de Ricqlès A (2009) L'origine et l'évolution des oiseaux: 35 années du progrès. *Comptes Rendus Pal Evol* 8:257–280
- Paul GS (2002) *Dinosaurs of the air: the evolution and loss of flight in dinosaurs and birds*. Johns Hopkins University Press, Baltimore
- Perez-Moreno BP, Sanz JL, Buscalioni AD, Moratalla JJ, Ortega F, Raskin-Gutman D (1994) A unique multitoothed ornithomimosaur from the Lower Cretaceous of Spain. *Nature* 370:363–367
- Perle A, Norell MA, Chiappe LM, Clark JM (1993) Flightless bird from the Cretaceous of Mongolia. *Nature* 362:623–626
- Prum RO (2002) Why ornithologists should care about the theropod origin of birds. *Auk* 119:1–17
- Senter P, Robins JH (2003) Taxonomic status of the specimens of *Archaeopteryx*. *J Vertebr Paleontol* 23:961–965
- Sereno PC (1999) The evolution of dinosaurs. *Science* 284:2137–2147
- Sereno PC (2001) Alvarezsaurids: Birds or ornithomimosaur? In: Gauthier J, Gall LF (eds) *New perspectives on the origin and early evolution of birds*. Yale Peabody Museum, New Haven, pp 363–385
- Sereno PC, Dutheil DB, Iarochene M, Larsson HCE, Lyon GH, Magwene PM, Sidor CA, Varricchio DJ, Wilson JA (1996) Predatory dinosaurs from the Sahara and the Late Cretaceous faunal differentiation. *Science* 272:986–991
- Sereno PC, Martinez RN, Wilson JA, Varricchio DJ, Alcober OA, Larsson HCE (2008) Evidence for avian intrathoracic air sacs in a new predatory dinosaur from Argentina. *PLoS ONE* 3:e3303
- Therrien F, Henderson DM (2007) My theropod is bigger than yours ... or not: estimating body size from skull length in theropods. *J Vertebr Paleontol* 27:108–115

- Van Tuinen M (2009) Birds (Aves). In: Blair Hedges S, Kumar S (eds) The timetree of life. Oxford University Press, New York, pp 409–411
- Vargas AO, Fallon JF (2005) Birds have Dinosaur wings: the molecular evidence. *J Exp Zool* 304:86–90
- Varricchio DV, Jackson F (2004) Two eggs sunny-side up: reproductive physiology in the dinosaur *Troodon formosus*. In: Currie PJ, Koppelhus EB, Shugar MA, Wright JL (eds) Feathered dragons: studies on the transition from dinosaurs to birds. Indiana University Press, Bloomington, pp 215–233
- Wagner A (1861) Neue Beiträge zur Kenntnis der urweltlichen Fauna des lithographischen Schiefers. *Abh Bayer Akad Wiss* 2:67–124
- Wallace AR (1858) On the tendency of varieties to depart indefinitely from the original type. *J Proc Linn Soc Lond* 3:45–62
- Wallace AR (1864) On the parrots of the Malayan Region with remarks on their habits, distribution, and affinities, and the descriptions of two new species. *Proc Zool Soc Lond* 1864:272–295
- Wallace AR (1865) Description of new birds from the Malay Archipelago. *Proc Zool Soc Lond* 1865:474–481
- Wallace AR (1868) On the raptorial birds of the Malay Archipelago. *Ibis* (ns) 4:1–27
- Wallace AR (1869) The Malay Archipelago: the land of the Orang-Utan and the bird of paradise. A narrative of travel with studies of man and nature, Vols. 1, 2. MacMillan & Co, London
- Wallace AR (1889) Darwinism: an exposition of the theory of natural selection with some of its applications. MacMillan & Co, London/New York
- Wallace AR (1910) The world of life: a manifestation of creative power, directive mind and ultimate purpose. Chapman and Hall, London
- Witmer LM (2002) The debate on avian ancestry: phylogeny, function, and fossils. In: Chiappe LM, Witmer LM (eds) Mesozoic birds: above the heads of dinosaurs. University of California Press, Berkeley, pp 3–30
- Xu X, Norell M (2004) A new troodontid dinosaur from China with avian-like sleeping posture. *Nature* 431:838–841
- Xu X, Zhou Z, Wang X, Kuang X, Zhang F, Du X (2003) Four-winged dinosaurs from China. *Nature* 421:335–340
- Xu X, You H, Du K, Han F (2011) An *Archaeopteryx*-like theropod from China and the origin of Avialae. *Nature* 475:465–470
- Zheng X, Zhong Z, Xiaoli W, Fucheng Z, Xiaomei Z, Yan W, Guangjin W, Shuo W, Xing X (2013) Hind wings in basal birds and the evolution of leg feathers. *Science* 339:1309–1312
- Zhou Z, Clarke JA, Zhang F (2002) *Archaeoraptor's* better half. *Nature* 420:285